I N S T I T U T D E
S T A T I S T I Q U E

UNIVERSITÉ CATHOLIQUE DE LOUVAIN

D I S C U S S I O N
P A P E R

0302

ANALYSIS OF BIVARIATE TAIL DEPENDENCE USING EXTREME VALUE COPULAS: AN APPLICATION TO THE SOA MEDICAL LARGE CLAIMS DATABASE

A.C. CEBRIÁN, M. DENUIT and P. LAMBERT

http://www.stat.ucl.ac.be

Analysis of bivariate tail dependence using extreme value copulas: an application to the SOA medical large claims database

Ana C. Cebrián*, Michel Denuit†‡ and Philippe Lambert†
*Dpto. Metodos Estadisticos. Ed. Matematicas
Universidad de Zaragoza
Zaragoza 50009 Spain

†Institut de Statistique
Université Catholique de Louvain
B-1348 Louvain-la-Neuve, Belgium

‡Institut des Sciences Actuarielles
Université Catholique de Louvain
B-1348 Louvain-la-Neuve, Belgium

January 23, 2003

Abstract

The aim of this work is to analyze the dependence structure between losses and ALAE's relating to large claims using extreme value copulas. We propose a procedure to select and estimate the copula based on a parametric estimation of the dependence function. An application to the evaluation of reinsurance premiums is performed in group medical insurance. It clearly enhances the relevance of the approach.

Key words and phrases: extreme value copulas, bivariate extreme value distributions, dependence function, excess-of-loss reinsurance premiums.

1 Introduction and Motivation

1.1 The situation under study

Quite often, actuaries face processes involving pairs of correlated variables. A fine example is the classical Loss-ALAE problem, examined e.g. by Frees and Valdez (1998) and Klugman and Parsa (1999). Here ALAE's are type of insurance company expenses that are specifically attributable to the settlement of individual claims such as lawyers' fees and claims investigation expenses. In order to price an excess-of-loss reinsurance treaty when the reinsurer shares the claim settlement costs, the dependence between losses and ALAE's has to be accounted for.

In this work we are going to deal with a database about medical insurance large claims, available from the website of the Society of Actuaries (`http://www.soa.org`), where the total claim corresponds to the sum of two amounts, the "Hospital" (Loss) and "Other" (ALAE) expenses. Since the random variables considered are continuous, we propose an approach based on copulas to describe the joint distribution of the pair (Loss,ALAE).

1.2 Data description

The dataset we are going to work with is part of a larger database that records medical claim amounts exceeding \$25,000. The study by Grazier *et al.* (1997) collects information from 26 insurers. The 171,000 claims recorded are part of a database including about 3,000,000 claims over the years 1991-92. The total amount paid by the insurer for each claim (coded as Total), which behaviour was studied in Cebrián *et al.* (2001), is the sum of hospital charges (Loss) and other expenses (ALAE). The database only records Total amounts exceeding \$25,000.

	Mean	Q_{25}	Q_{50}	Q_{75}	Min.	Max.	St. dev.
Loss	63,450	31,360	42,310	66,130	25,000	1,430,000	74,665
ALAE	18,920	8,088	13,940	22,850	23	568,000	22,073
Total	82,370	43,040	58,560	88,500	25,320	1,530,000	85,523

Table 1.1: *Claims summary statistics.*

In this study only the 1991 record relating to Plan type 4 is used. Since

the main interest is the analysis of extreme claims, only observations with Hospital charges greater than 25,000$ are considered. The size of this sample is 3,901. Univariate descriptive statistics for the the three variables are provided in Table 1.1.

The relationship between Loss and ALAE is illustrated in Figure 1, where the scatterplots of the two variables, in original and log-scale are shown. Although there is a clear dependence, it is not simple to model. Tests for independence (Pearson with $\widehat{r} = 0.380$, Spearman with $\widehat{\rho} = 0.437$ and Kendall with $\widehat{\tau} = 0.305$) all clearly reject independence (p-values less than 10^{-2}).



Figure 1: *Bivariate plots of Loss and ALAE on original and log-scale.*

1.3 Aim of the study

Our purpose is to price an excess-of-loss reinsurance treaty where the settlement costs are shared between the reinsurer and insurer on a proportional basis. Specifically, given a realization of (Loss,ALAE), the reinsurance indemnity amounts to

$$g(Loss, ALAE) = \begin{cases} 0 & \text{if } Loss \leq R, \\ Loss - R + \frac{Loss-R}{Loss}\ ALAE & \text{if } Loss > R. \end{cases} \tag{1.1}$$

The pure premium relating to this reinsurance treaty is expressed in terms of

$$\pi = E[g(Loss, ALAE)].$$

The pure reinsurance premium is obtained by multiplying π by the expected number of claims generated by the portfolio. Clearly, the joint distribution of the pair (Loss,ALAE) has to be modelled to compute π. This is precisely the aim of the next section.

2 Modelling the dependence between Loss and ALAE

2.1 Copulas

2.1.1 Sklar's construction

We briefly recall hereafter the basic features of the copula construction. We refer the interested reader to Frees and Valdez (1998) for an introduction to copulas with applications in actuarial science.

A copula is a distribution function $C : [0,1]^2 \to [0,1]$ with unit uniform marginals. This function entirely describes the dependence structure between two random variables X_1 and X_2. The interest of this statistical tool originates from the following result, referred to as Sklar's theorem: Given a bidimensional distribution function F, with univariate margins F_1 and F_2, there exists a copula C such that for all $(x_1, x_2) \in \mathbb{R}^2$,

$$F(x_1, x_2) = C(F_1(x_1), F_2(x_2)). \tag{2.1}$$

Conversely, if C is a copula and F_1 and F_2 are distribution functions then the function F defined by (2.1) is a bidimensional distribution with margins

F_1 and F_2. Moreover, if F_1 and F_2 are both continuous, then C is uniquely defined.

Henceforth, we assume that the marginals F_1 and F_2 are both continuous. Then, for any $(u_1, u_2) \in [0,1]^2$,

$$\begin{aligned} C(u_1, u_2) &= P[F_1(X_1) \leq u_1, F_2(X_2) \leq u_2] \\ &= P[X_1 \leq F_1^{-1}(u_1), X_1 \leq F_2^{-1}(u_2)] \\ &= F\left(F_1^{-1}(u_1), F_2^{-1}(u_2)\right) \end{aligned} \tag{2.2}$$

where the F_i^{-1}'s, $i = 1, 2$, are the quantile functions associated with the F_i's, i.e.

$$F_i^{-1}(p) = \inf\{x \in \mathbb{R} | F_i(x) \geq p\}, \;\; p \in (0,1).$$

As it can be seen from (2.2) the copula can be recovered from the knowledge of the joint distribution F and of its margins F_j, $j = 1, 2$.

2.1.2 Copulas and dependence measures

Copulas capture the dependence between two variables, regardless of the scale in which each variable is measured. More precisely, given two arbitrary functions g_1 and g_2 strictly increasing over the range of X_1 and X_2, then the transformed variables $g_1(X_1)$ and $g_2(X_2)$ have the same copula than X_1 and X_2.

Moreover, usual measures of correlation between two variables, such as the Spearman's and Kendall's coefficients can be expressed uniquely in terms of their copula. Specifically, the following representations hold true:

$$\begin{aligned} \rho &= 12 \int_0^1 \int_0^1 \{C(u,v) - uv\} du dv \\ \tau &= 4 \int_0^1 \int_0^1 C(u,v) dC(u,v) - 1. \end{aligned}$$

2.2 Extreme value copulas

In the situation of interest in this paper, the combination of the copula construction with extreme value theory seems promising. This is precisely the aim of this section.

2.2.1 Definition

A random vector (X_1, X_2) is said to conform to a Bivariate Extreme Value Distribution (BEVD, in short) with unit exponential margins if,

$$P[X_1 > y_1] = \exp(-y_1) \text{ and } P[X_2 > y_2] = \exp(-y_2)$$

for $y_1, y_2 > 0$ and the joint survival function $\bar{F}$ verifies,

$$\bar{F}^n(y_1, y_2) = \bar{F}(ny_1, ny_2)$$

for any integer $n \geq 1$. For more details, see e.g. Tawn (1988). A bivariate vector follows such a distribution if and only if, the survival function can be expressed as

$$\bar{F}(y_1, y_2) = \exp\left[-(y_1 + y_2)A\left(\frac{y_2}{y_1 + y_2}\right)\right]$$

for any $y_1, y_2 > 0$, where

$$A(w) = \int_0^1 \max[(1-w)q, w(1-q)]dL(q)$$

for some positive finite measure L on $[0, 1]$.

Expressing this relationship in terms of uniform margins, we obtain

$$C(u_1, u_2) = \exp\left\{[\log(u_1) + \log(u_2)]A\left[\frac{\log(u_2)}{\log(u_1) + \log(u_2)}\right]\right\}$$

for $0 \leq u_1, u_2 \leq 1$. This function is known as the *bivariate extreme value copula* and it allows to model dependence between the components of a bivariate vector of variables which represent the largest values of two characteristics observed over the same period of time.

2.2.2 Dependence function

The function A is called the *dependence function* of (X_1, X_2) and it satisfies some properties:

- $A(0) = A(1) = 1$.
- $\max(w, 1-w) \leq A(w) \leq 1$ for $0 \leq w \leq 1$. Moreover, if $A(w) = 1$ then (X_1, X_2) are independent and if $A(w) = \max(w, 1-w)$, then (X_1, X_2) are perfectly dependent (or comonotonic).
- $A(w)$ is a convex function in the region $0 \leq w \leq 1$.

2.3 Some examples

2.3.1 Logistic model

The dependence function is

$$A(w) = [(1-w)^r + w^r]^{1/r}$$

with $r \geq 1$. The corresponding copula function is given by

$$C(u_1, u_2) = \exp\left\{-[(-\ln u_1)^r + (-\ln u_2)^r]^{1/r}\right\}.$$

Independence and complete dependence correspond to $r = 1$ and $r = \infty$ respectively. In this model the variables are exchangeable.

2.3.2 Asymmetric logistic model

The dependence function is

$$A(w) = [\theta^r(1-w)^r + \phi^r w^r]^{1/r} + (\theta - \phi)w + 1 - \theta$$

with $\theta \geq 0$, $\phi \leq 1$, and $r \geq 1$ and the copula function

$$C(u_1, u_2) = \exp\left\{(1-\theta)\ln u_1 + (1-\phi)\ln u_2 - [(-\ln u_1)^r\theta^r + (-\ln u_2)^r\phi^r]^{1/r}\right\}.$$

This model is very flexible and contains several of the existing models such as the logistic ($\phi = \theta = 1$), the biextremal, the dual of the biextremal, the Gumbel, as well as a mixture of logistic and independence models. Complete dependence corresponds to $\theta = \phi = 1$ and $r = \infty$ whereas independence corresponds to $\theta = 0$ or $\phi = 0$ or $r = 1$.

2.3.3 Mixed model

The dependence function is

$$A(w) = \theta w^2 - \theta w + 1$$

and the corresponding copula function is

$$C(u_1, u_2) = u_1 u_2 \exp\left\{-\theta \frac{\ln u_1 \ln u_2}{\ln(u_1 u_2)}\right\}$$

with $0 \leq \theta \leq 1$. Independence corresponds to $\theta = 0$ but we cannot have complete dependence. In this model the variables are exchangeable.

2.4 Kendall's tau

Some important properties of the variables can be formulated in terms of this function A. In particular Kendall's τ is given by

$$\tau = \int_0^1 \frac{t(1-t)}{A(t)} A''(t)dt. \tag{2.3}$$

For the models presented above, τ_A is given by

1. Logistic model: $\tau_A = 1 - 1/r$.
2. Mixed model: $\tau_A = \frac{8 \arctan \sqrt{\theta/(4-\theta)}}{\sqrt{\theta(4-\theta)}} - 2$.
3. Asymmetric Logistic model: $\tau_A = \int_0^1 \frac{w(1-w)(r-1)\theta^r \phi^r (w-w^2)^{r-2}}{[\theta^r(1-w)^r + \phi^r w^r]^2} dw$

For the asymmetric logistic model, a closed expression for τ_A does not exist. Therefore, one needs to resort to numerical methods to evaluate τ_A.

3 Estimation of the copula C and its probability integral transform K

3.1 Copula probability integral transform

Genest and Rivest (2000) suggest a general formula for computing the distribution function K of the random variable

$$V = F(X_1, X_2) = C(F_1(X_1), F_2(X_2)).$$

In contrast to the univariate case, it is not true that V is uniformly distributed over $[0, 1]$. In fact, it is well-known that any copula is constrained from below and above by the Fréchet bounds as

$$\max(0, u_1 + u_2 - 1) \leq C(u_1, u_2) \leq \min(u_1, u_2)$$

holds for $0 \leq u_1, u_2 \leq 1$. Consequently, any distribution function K for $F(X_1, X_2)$ has to fulfill the inequalities

$$v \leq K(v) \leq 1$$

for $0 \leq v \leq 1$. Moreover, $K(v) = v$ and $K(v) = 1$ correspond to the two extreme cases where the random variables X_1 and X_2 are in perfect positive dependence (comonotonicity) and in perfect negative dependence (countermonotonicity), respectively.

It is not possible to characterize F from the knowledge of K alone since it does not contain any information about the marginal distributions of X_1 and X_2. However, K contains useful information about the dependence structure underlying F. It can be proved that

$$K(v) = v - \lambda(v)$$

with

$$\lambda(v) = -\int_v^1 \frac{\partial C(x_1, x_2^{x_1 v})}{\partial x_1} dx_1$$

and $x_2^{x_1 v}$ the root of the equation

$$C(x_1, x_2^{x_1 v}) = v.$$

3.2 Computing K for bivariate extreme value copulas

For the extreme value copulas discussed in Section 2.2, we obtain

$$\lambda(v) = -(1 - \tau_A) v \log(v)$$

where τ_A is given by (2.3) so that

$$K(v) = v - (1 - \tau_A) v \log(v). \tag{3.1}$$

Obviously, in order to estimate K via (3.1), only τ_A needs to be estimated from the data.

3.3 Parameter estimation

Given a random sample of size n, $(X_{11}, X_{21}), \ldots (X_{1n}, X_{2n})$ say with copula function C describing the dependence between each pair of variables and marginal density functions f_1 and f_2, the corresponding joint probability density function is

$$f(x_1, x_2) = f_1(x_1) f_2(x_2) C''(F_1(x_1), F_2(x_2); p)$$

where p is a vector of parameters and

$$C''(u_1, u_2; p) = \frac{\partial^2}{\partial u_1 \partial u_2} C(u_1, u_2; p).$$

Assuming known margins, only C'' depends on the parameters to be estimated and the function

$$L(p) = \sum_{i=1}^{n} \log C''(u_{1i}, u_{2i}; p) \text{ with } u_{ji} = F_j^{-1}(x_{ji}), \;\; j = 1, 2, \tag{3.2}$$

must be maximized, using for example algorithms based on quasi-Newton methods.

3.4 Nonparametric estimation of C

To check the validity of the parametric models, it would be desirable to compare them with nonparametric estimators. In particular, we make use of the one proposed by Scaillet (2000). This author developed a kernel based approach without putting any particular parametric a priori on the dependence structure between margins.

To estimate the copula function at given points, $\{u_i = (u_{1i}, u_{2i})\}_{i=1,\ldots,d}$, with $u_{ji} \in (0, 1)$, a simple plug-in method in expression (2.2) is used, resulting in,

$$\hat{C}(u_1, u_2) = \hat{F}(\hat{F}_1^{-1}(u_1), \hat{F}_2^{-1}(u_2))$$

where $\widehat{F}$, $\widehat{F}_1$ and $\widehat{F}_2$ represent estimators of the bivariate distribution function F and of the corresponding marginal distributions F_1 and F_2. In this work, the empirical distribution function will be used to estimate each of the marginal functions F_1 and F_2, but kernel or parametric estimators (if proper distributions are found to fit the margins) could also be used.

3.5 Nonparametric estimation of K

In order to compare the performed estimations, it would be also desirable to have available an estimator of K. One advantage of this function is that, since it is unidimensional, it can be easily represented using usual graphical tools. A kernel estimator, based on the pseudo-observations

$$c_i = \hat{C}(\hat{F}(x_{1i}), \hat{F}(x_{2i})),$$

will be used to estimate the function K.

Model	Parameters	$\widehat{\tau_A}$	(3.2)	AIC
Logistic	$\widehat{r}$ =1.406	0.289	-7,321.3	14,643
Asym. Log.	$\widehat{\theta}$ =0.983, $\widehat{\phi}$ =0.825, $\widehat{r}$ =1.486	0.288	-7,315.7	14,631
Mixed	$\widehat{\theta}$ =0.738	0.289	-7,327.3	14,654

Table 4.1: *Summary of dependence function estimation.*

4 Loss-ALAE analysis

4.1 Estimation results

Table 4.1 summarizes estimations for the three parametric models discussed in Section 2.3. Not only the value of the function (3.2) is given, but also an AIC measure,

$$AIC = -2L + 2n_p/n$$

where L has been defined in (3.2) and n_p is the number of parameters, is calculated as goodness of fit measures for each model.

In Figure 2 the three estimations of the dependence funtion are represented. It can be seen that the three models provide similar results and it seems all of them are able to represent the dependence structure. Also the estimated Kendall τ is almost the same for the three models.

In plots of Figure 3 we represent for each model the differences between the nonparametric and the constrained estimations of the copula function. From the upper left graph, the difference between the nonparametric estimation of the copula and its independent counterpart is always positive, indicating a positive dependence between Loss and ALAE. The results from all the parametric models are rather similar, but the mixed model looks slightly better than the others (since the differences are smaller for this model).

In Figure 4 we compare the parametric estimations of K with a kernel estimation of this function calculated from the copula sample $\widehat{c}_i = \hat{C}(\hat{F}_1(x_{1i}), \hat{F}_2(x_{2i}))$. Again the independent model is rather bad, producing large departures from the nonparametric estimation. No significant differences are observed between the three parametric models.

A final possibility to check the model is to compare the quantiles of the parametric and the nonparametric estimation of the copula using qq-plots. The plots corresponding to the independent and the parametric models are



Figure 2: *Parametric estimations of the dependence function A.*

shown in Figure 5. Again, independence excepted, the three parametric models behave well on the data set under study.

4.2 Goodness of fit tests

In addition to the graphical tools, some more objective measures and tests are desirable to complete the goodness of fit analysis. First, we performed a general test proposed by Ghoudi et al. (1998) to verify whether the copula belongs to the bivariate extreme value family without specifying a particular dependence function. It is based on the distribution function of the copula, K; more precisely, the hypothesis $H_0 : K(v) = K_\tau(v)$ with K_τ of the form (3.1) for some $\tau \in (0, 1)$ is checked using the test statistic

$$\begin{aligned} S_n &= \frac{8}{n(n-1)} \sum_{i \neq j}^{n} \delta[(X_{1i}, x_{2i}), (X_{1j}, x_{2j})] - \\ &\quad \frac{9}{n(n-1)(n-2)} \sum_{i \neq j \neq k}^{n} \delta[(X_{1i}, x_{2i}), (X_{1j}, x_{2j})] \delta[(X_{1i}, x_{2i}), (X_{1k}, x_{2k})] - 1, \end{aligned}$$



Figure 3: *Differences between the nonparametric and the parametric estimation of the copula.*

where

$$\delta[(x_1, y_1), (x_2, y_2)] = \begin{cases} 1 \text{ if } x_1 \geq x_2 \text{ and } y_1 \geq y_2 \\ 0 \text{ otherwise.} \end{cases}$$

The standarized test statistic $|S_n|/\hat{Var}(S_n)$ is asymptotically normal and a jackknife method is suggested to estimate the variance,

$$\hat{Var}(S_n) = \frac{n-1}{n} \sum_{i=1}^{n} \left[S_n^{-i} - S_n \right]^2 .$$

Satisfactory results were found with an observed value of S_n equal to 1.016 and the corresponding p-value equal to 0.31.

After that, we checked the goodness of fit of the different dependence functions. We used a Kolmogorov-Smirnov type test to compare the empirical function distribution of the sample $\hat{C}(\hat{F}_1(x_{1i}), \hat{F}_2(x_{2i}))$ to the estimation of the function K estimated under one of the parametric models presented in Section 2.3 and evaluated in the same observations. As we can see in Table 4.2, according to the K-S test no model is rejected with any usual confidence level.

Test	Logistic	Asym. Logistic.	Mixed
K-S test statistic	0.0121	0.0118	0.0118
p-v	0.61	0.64	0.64

Table 4.2: Goodness of fit tests.

5 Reinsurance premiums

Let us now analyze the impact of the dependence structure on premium valuation in reinsurance treaties. We consider a reinsurance treaty on a policy with unlimited liability and insurer's retention R. Assuming a pro-rata sharing of expenses, the reinsurer's payment for a given realization of (Loss,ALAE) is described by the function (1.1).

As it is demonstrated in Joe (1997), any extreme value copula verifies the association property. This means that if g_1 and $g_2 : \mathbb{R}^2 \to \mathbb{R}$ are non-decreasing functions, the inequality

$$E[g_1(X_1, X_2)g_2(X_1, X_2)] \geq E[g_1(X_1, X_2)]E[g_2(X_1, X_2)]$$

holds true. Association is a rather strong dependence property. In particular it implies positive quadrant dependence, that is

$$P(X_1 > x_1, X_2 > x_2) \geq P(X_1 > x_1)P(X_2 > x_2).$$

From Denuit, Dhaene and Ribas (2001), it is known that the premium π computed under the independence hypothesis for (Loss,ALAE) will be smaller than the premium computed under an extreme value copula.

The results in Table 5.1 provide the premiums the reinsurer would have assessed to cover Loss and ALAE charges according to various insurer's retention. Three situations have been considered:

1. Assuming independence, an estimator of the premium π is

$$\widehat{\pi} = \frac{1}{n^2} \sum_{t=1}^{n} \sum_{t'=1}^{n} g(Loss_t, ALAE_{t'}).$$

   Since g is supermodular, this value is a lower bound to the unknown π under positive quadrant dependence for Loss and ALAE. It is worth to mention that Figure 3 (upper left panel) precisely suggested such a positive dependence).

2. Applying the classical comonotonic approximation for (Loss,ALAE) we obtain,

$$\widehat{\pi} = \frac{1}{n} \sum_{t=1}^{n} g\left(Loss_t, \widehat{F}_{ALAE}^{-1}\left[\widehat{F}_{Loss}(Loss_t)\right]\right).$$

   Because of the supermodularity of g, this value is an upper bound π.

3. If we consider the dependence structure, we have two possibilities,

   a.- either we do not specify any model for the dependence, i.e. we compute the empirical analogue of π given by

$$\widehat{\pi} = \frac{1}{n} \sum_{t=1}^{n} g(Loss_t, ALAE_t);$$

   b.- or we estimate π under one of the copula models presented in Section 2.3. Under these models, the pure premium can be estimated by

$$\hat{\pi} = \hat{E}[g(Loss, ALAE)] = \int_{25,000}^{\infty} \int_{0}^{\infty} g(l, a) \hat{f}_{Loss,ALAE}(l, a)\, dl\, da;$$

where

$$\widehat{f}_{Loss,ALAE}(l,a) = \widehat{f}_{Loss}(l)\widehat{f}_{ALAE}(a)\widehat{C''}[\widehat{F}_{Loss}(l), \widehat{F}_{ALAE}(a)].$$

Since no parametric models have been posed for the margins, the marginal densities are estimated using kernel methods. Having an estimator of the bivariate density function available, the double integral is then calculated using numerical procedures. Since the integrand is a rather complicated function, a powerful software to perform analytical calculations is necessary; in this case MathCad 2000 has been used. Although the procedure is not complicated, it is rather time consuming, mainly due to the evaluations of the kernel estimators of the density and distribution functions.

The $\hat{\pi}$ values for $R = 25,000,\ 50,000,\ 100,000,\ 500,000,\ 1,000,000$ estimated under the different models are shown in Table 5.1. We see that substantial mispricing could result from the independence hypothesis, while the comononotic approximation is too conservative. Thus, as expected, independence generates lower premiums than those taking into account dependence of the data, which themselves are smaller than those based on the comonotonic assumption. Two estimators considering dependence are shown, the one based on the empiric dependence of the data and the one based on the extreme value copula with the mixed model, that provided the best fit to the bivariate sample. Comparing both estimators, we observe that they are rather similar for small R values, but higher the values of R, higher the relative differences between them. It must be remembered that, in fact, the main drawback of nonparametric estimation for π is that for high R values the sample size is not large enough to get reliable estimations.

6 Conclusions

To conclude we can say that the results show the importance to consider the dependence structure of the two variables: substantial mispricing can result from the independence assumption, while the comonotonic approximation seriously overestimates the premium.

Given the characteristics of the analyzed variables, the use of bivariate extreme value copulas is intellectually satisfying. Moreover, the SOA medical

R	25,000	50,000	100,000	500,000	1,000,000
Independence	46,365.7	25,642.3	12,785.5	1,247.1	230.3
empirical	48,350.6	27,639.1	13,842.3	1,336.6	244.7
(# obs. Loss$> R$)	(3901)	(1516)	(470)	(17)	(3)
Mixed model	48,357.5	27,804.6	14,188.3	1,431.6	269.3
Comonotonicity	50,318.8	29,760.7	15,374.9	1,570.6	302.1

Table 5.1: Pure premiums for a reinsurance treaty with retention R.

large claims database supported these models. Therefore, extreme value copulas should be considered for bivariate reinsurance pricing in the future.

Acknowledgements

Ana Cebrián thanks the Université catholique de Louvain, Louvain-la-Neuve, Belgium, for financial support through a FSR research grant.

Financial support from the contract 'Projet d'Actions de Recherche Concertées' nr 98/03-217 from the Belgian government, and from the IAP research network nr P5/24 of the Belgian State (Federal Office for Scientific, Technical and Cultural Affairs) is gratefully acknowledged.

References

[1] Cebrian, A., Denuit, M., and Lambert, Ph. (2001). Determination of the PML: a case study in group medical insurance. Discussion Paper 01-03, Institut de Statistique, Universit Catholique de Louvain, Louvain-la-Neuve, Belgium.

[2] Denuit, M., Dhaene, J., and Ribas, C. (2001). Does positive dependence between individual risks increase stop-loss premiums? *Insurance: Mathematics and Economics* **28**, 305-308.

[3] Frees, E.W. and Valdez, E. A. (1998). Understanding relationships using copulas. *North American Actuarial Journal* **2**, 1-15.

[4] Genest, C. and Rivest, L.P. (2000). On the Multivariate Probability Integral Transformation. *Statistics and Probability Letters* **53**, 391-399.

[5] Ghoudi, K., Khoudraji, A. and Rivest L.P. (1998). Propriétés Statistiques des Copules de Valeurs Extrêmes Bidimensionnelles. *The Canadian Journal of Statistics* **26** No 1, 187-197.

[6] Grazier, K.L., and G'Sell Associates (1997). *Group Medical Insurance Large Claims Database and Collection*. SOA Monograph M-HB97-1.

[7] Joe, H. (1997). *Multivariate Models and Dependence Concepts*. Chapman & Hall, London

[8] Klugman, S. A. and Parsa, R. (1999). Fitting Bivariate Loss Distributions with Copulas. *Insrance: Mathematics and Economics* **24**, 139-148.

[9] Tawn, J. (1988). Bivariate Extreme Value Theory: Models and Estimation. *Biometrika* **75**, 397-415.

[10] Scaillet, O. (2000). Nonparametric Estimation of Copulas for Time Series. Downloadable on http://www.hec.unige.ch.



Figure 4: *Estimation of the distribution function K of the copula.*



Figure 5: *QQ-plots of the copula quantiles.*