

I N S T I T U T D E
S T A T I S T I Q U E

UNIVERSITÉ CATHOLIQUE DE LOUVAIN



D I S C U S S I O N
P A P E R

0912

**GOODNESS-OF-FIT TESTS FOR THE ERROR
DISTRIBUTION IN NONPARAMETRIC
REGRESSION**

HEUCHENNE, C. and I. VAN KEILEGOM

This file can be downloaded from
<http://www.stat.ucl.ac.be/ISpub>

Goodness-of-fit tests for the error distribution in nonparametric regression

Cédric HEUCHENNE *

HEC-Management School

University of Liège

Ingrid VAN KEILEGOM †

Institute of Statistics

Université catholique de Louvain

June 15, 2009

Abstract

Suppose the random vector (X, Y) satisfies the regression model $Y = m(X) + \sigma(X)\varepsilon$, where $m(\cdot) = E(Y|\cdot)$, $\sigma^2(\cdot) = \text{Var}(Y|\cdot)$ and ε is independent of X . The covariate X is d -dimensional ($d \geq 1$), the response Y is one-dimensional, and m and σ are unknown but smooth functions. In this paper we study goodness-of-fit tests for the parametric form of the error distribution under this model, without assuming any parametric form for m or σ . The proposed tests are based on the difference between a nonparametric estimator of the error distribution and an estimator obtained under the null hypothesis of a parametric model. The large sample properties of the proposed test statistics are obtained, as well as those of the estimator of the parameter vector under the null hypothesis. Finally, the finite sample behavior of the proposed statistics, and the selection of the bandwidths for estimating m and σ are extensively studied via simulations.

Key Words: Bandwidth selection; bootstrap; error distribution; goodness-of-fit tests; least squares estimation; local polynomial estimation; maximum likelihood; nonparametric regression.

*Financial support from IAP research network P6/03 of the Belgian Government (Belgian Science Policy) is gratefully acknowledged.

†Financial support from IAP research network P6/03 of the Belgian Government (Belgian Science Policy), and from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013) / ERC Grant agreement No. 203650 is gratefully acknowledged.

1 Introduction

Suppose the d -dimensional random vector X and the random variable Y satisfy the following heteroscedastic regression model :

$$Y = m(X) + \sigma(X)\varepsilon, \quad (1.1)$$

where ε is independent of the random vector X , $m(X) = E[Y|X]$ and $\sigma^2(X) = \text{Var}[Y|X]$.

In the literature, many papers have been devoted to testing the form of $m(\cdot)$ and $\sigma(\cdot)$. Goodness-of-fit tests for $m(\cdot)$ were studied by e.g. Härdle and Mammen (1993), Stute (1997), Dette and Munk (1998), Alcalá, Cristóbal and González Manteiga (1999), Dette (1999), Fan, Zhang and Zhang (2001) and Van Keilegom, González Manteiga and Sánchez Sellero (2008), among many others. Testing parametric models for $\sigma(\cdot)$ has also been investigated, see e.g., among others, Dette, Neumeyer and Van Keilegom (2007). In the above papers the form of the error distribution is left unspecified.

In this paper, we are interested in testing hypotheses concerning the form of this error distribution, without making any assumption regarding the form of the regression function $m(\cdot)$ and the variance function $\sigma^2(\cdot)$, except for smoothness assumptions. Knowing that ε follows a certain parametric distribution offers important advantages when doing inference for the functions $m(\cdot)$ and $\sigma^2(\cdot)$.

Consider the parametric class $\mathcal{F} = \{F_{\varepsilon\theta} : \theta \in \Theta\}$ of distribution functions, where Θ is a compact subset of \mathbb{R}^k and k is a positive integer. We denote the true value of θ by θ_0 , and the distribution of the error ε by $F_\varepsilon(y) = P(\varepsilon \leq y)$. The aim of this paper is to test the hypothesis

$$H_0 : F_\varepsilon \in \mathcal{F} \text{ versus } H_1 : F_\varepsilon \notin \mathcal{F}. \quad (1.2)$$

This testing problem has been studied by Jiménez Gamero, Muñoz García and Pino Mejías (2005) for linear regression models and by Neumeyer, Dette and Nagel (2006) for linear and nonparametric regression models. In the latter paper, the authors work with a generic estimator for θ_0 that satisfies a certain asymptotic representation, and they obtain the asymptotic theory for their proposed test statistic under the assumption that such an estimator of θ_0 exists (see their assumption B5). In this paper, we work with a specific estimator for θ_0 , obtained by using a maximum likelihood approach, and we work out in detail the asymptotic properties of this estimator, and of the proposed test statistics. These test statistics are based on a Kolmogorov-Smirnov and a Cramér-von Mises distance between an estimator of the error distribution obtained under H_0 and a

completely nonparametric estimator. Under the null hypothesis H_0 , we show that the estimator of θ_0 and the test statistics reach the same rate of convergence as in the usual case where $m(\cdot)$ and $\sigma(\cdot)$ are parametric functions. The asymptotic results are largely based on the work of Neumeyer and Van Keilegom (2009) regarding the estimation of the error distribution under model (1.1), and on the results obtained by Chen, Linton and Van Keilegom (2003) regarding inference for general semiparametric models involving non-smooth criterion functions.

In practice, the power of the tests are somewhat sensitive to the choice of the bandwidths used to estimate $m(\cdot)$ and $\sigma(\cdot)$. We therefore study six different procedures to select these bandwidths. As it turns out, the power seems to be higher when loss functions using (a cross validation version of) the residuals themselves are involved.

The paper is organized as follows. In the next section, the estimator of θ_0 and the test statistic are described in detail. Section 3 summarizes the main asymptotic results including the asymptotic normality of the estimator of θ_0 and the weak convergence of the proposed test statistics under H_0 . In Section 4, we study the finite sample behavior of the test procedure and the selection of the smoothing parameters for $m(\cdot)$ and $\sigma(\cdot)$ through extended simulations, while the Appendix contains the assumptions and the proofs of the asymptotic results of Section 3.

2 Description of the method

Let $(X_1, Y_1), \dots, (X_n, Y_n)$ be an i.i.d. random sample generated from model (1.1), where the components of X_i are denoted by (X_{i1}, \dots, X_{id}) ($i = 1, \dots, n$). The distribution of ε and X are denoted by F_ε and F_X respectively, and their probability density functions by f_ε and f_X . We start by estimating the regression function $m(x)$ and the variance function $\sigma^2(x)$ at an arbitrary point $x = (x_1, \dots, x_d)$ in the support R_X of X , which we suppose to be a compact subset of \mathbb{R}^d . We estimate $m(x)$ by a local polynomial estimator of degree p , i.e. $\hat{m}(x) = \hat{\beta}_0$, where $\hat{\beta}_0$ is the first component of $\hat{\beta}$, which is the solution of the local minimization problem

$$\min_{\beta} \sum_{i=1}^n \{Y_i - P_i(\beta, x, p)\}^2 K_h(X_i - x), \quad (2.1)$$

where $P_i(\beta, x, p)$ is a polynomial of order p built up with all $0 \leq k \leq p$ products of factors of the form $X_{ij} - x_j$ ($j = 1, \dots, d$), and β is the vector of all coefficients of this polynomial. Here, for $u = (u_1, \dots, u_d) \in \mathbb{R}^d$, $K(u) = \prod_{j=1}^d k(u_j)$ is a d -dimensional product kernel,

k is a univariate kernel function, $h = (h_1, \dots, h_d)$ is a d -dimensional bandwidth vector converging to zero when n tends to infinity and $K_h(u) = \prod_{j=1}^d k(u_j/h_j)/h_j$. In the same way, $\hat{\sigma}^2(x) = \hat{\gamma}_0$ is the first component of $\hat{\gamma}$, which is the solution of the local minimization problem

$$\min_{\gamma} \sum_{i=1}^n \{(Y_i - \hat{m}_l(X_i))^2 - P_i(\gamma, x, q)\}^2 K_g(X_i - x), \quad (2.2)$$

where $\hat{m}_l(X_i)$, $i = 1, \dots, n$, is a local polynomial estimator obtained from (2.1) (except that h is replaced by l , which will be specified in Remark 2.1 below) and $P_i(\gamma, x, q)$, γ , $K_g(u)$ and $g = (g_1, \dots, g_d)$ are defined in a similar way as $P_i(\beta, x, p)$, β , $K_h(u)$ and h . An estimator for $\sigma(x)$, $\hat{\sigma}(x)$, will be simply obtained by taking the square root of $\hat{\gamma}_0$.

The nonparametric residuals can then be introduced into the likelihood function which will provide a vector of parameter estimators θ_n for θ_0 by solving the maximization problem

$$\max_{\theta \in \Theta} \sum_{i=1}^n \log f_{\varepsilon\theta}(\hat{\varepsilon}_i), \quad (2.3)$$

where $\hat{\varepsilon}_i = (Y_i - \hat{m}(X_i))/\hat{\sigma}(X_i)$ ($i = 1, \dots, n$).

Remark 2.1 (Choice of the smoothing parameters) The objective is to provide an easy and data-driven way to select the smoothing parameters in (2.1) and (2.2). To this end, we propose to compare six procedures in Section 4. Four are based on different least squares cross-validation ideas and two on maximum likelihood cross-validation ideas. More precisely, let

$$h_n = \operatorname{argmin}_h \sum_{j=1}^n (Y_j - \hat{m}_{h,-j}(X_j))^2, \quad (2.4)$$

$$g_{n1} = \operatorname{argmin}_g \sum_{j=1}^n \{(Y_j - \hat{m}_{g,-j}(X_j))^2 - \hat{\sigma}_{g,-j}^2(X_j)\}^2 \quad (2.5)$$

and

$$g_{n2} = \operatorname{argmin}_g \sum_{j=1}^n \{(Y_j - \hat{m}_{h_n}(X_j))^2 - \hat{\sigma}_{g,h,-j}^2(X_j)\}^2. \quad (2.6)$$

Here, $\hat{m}_{l,-j}(X_j)$ is a local polynomial estimator obtained by an expression of the type (2.1) for $x = X_j$ and h replaced by $l = g$ or h , but based on a sample for which the j^{th} data point has been removed. Moreover, $\hat{\sigma}_{g,-j}^2(X_j)$ is a local polynomial estimator

obtained from an expression of the type (2.2) for $x = X_j$, but based on the couples $(X_i, (Y_i - \hat{m}_{g,-i}(X_i))^2)$ ($i = 1, \dots, j-1, j+1, \dots, n$). Similarly, $\hat{\sigma}_{g,h,-j}^2(X_j)$ is obtained from the pairs $(X_i, (Y_i - \hat{m}_{h_n}(X_i))^2)$ ($i = 1, \dots, j-1, j+1, \dots, n$). The two first procedures that we will consider in detail in Section 4 are based on choosing h_n for h , g_{n1} (resp. h_n) for l and g_{n1} (resp. g_{n2}) for g .

An alternative idea for choosing h , g and l is to transform (2.3) in a joint maximization problem over θ , h , g and l . This could be simply achieved by jointly maximizing a cross-validation version of (2.3) with respect to θ , h , g and l . More precisely, we propose to obtain h_{ns} (for $h = g = l$) by solving

$$\max_{\theta, h} \sum_{j=1}^n \log f_{\varepsilon\theta}(\hat{\varepsilon}_{j,-j,h,s}) \quad (2.7)$$

for $s = 1, 2$, and where

$$\hat{\varepsilon}_{j,-j,h,s} = \frac{Y_j - \hat{m}_{h,-j}(X_j)}{\hat{\sigma}_{h,s}(X_j)} \quad (2.8)$$

for $\hat{\sigma}_{h,1}(X_j)$ a local polynomial estimator obtained from an expression of the type (2.2) for $x = X_j$ and $g = h$, but based on the couples $(X_i, (Y_i - \hat{m}_{h,-i}(X_i))^2)$ ($i = 1, \dots, n$) and $\hat{\sigma}_{h,2}(X_j) = \hat{\sigma}_{h,-j}(X_j)$.

Finally, an intermediate idea consists in directly using the residuals in cross-validation least squares minimization problems. That leads to choose h'_{ns} (for $h = g = l$) by solving

$$\min_h \sum_{i=1}^n \hat{\varepsilon}_{j,-j,h,s}^2 \quad (2.9)$$

for $s = 1, 2$.

To summarize, the six selection procedures that we propose for the bandwidths h and g are: (a) $(h, l, g) = (h_n, g_{n1}, g_{n1})$, (b) $(h, g) = (h_n, h_n, g_{n2})$, (c) $h = g = l = h_{n1}$, (d) $h = g = l = h_{n2}$, (e) $h = g = l = h'_{n1}$ and (f) $h = g = l = h'_{n2}$. Their practical performance will be studied in Section 4.

Remark 2.2 (Order of local polynomials) Apart from the practical choices for h , l and g discussed above, it is important to mention here the dependency between h (l and g) and the dimension d of X . Indeed, the sample size n should increase exponentially with d to preserve the convergence rates (curse of dimensionality). Consequently, for fixed sample size n , in order to compensate for this curse of dimensionality, the bandwidths h_j, l_j and g_j ($j = 1, \dots, d$) should increase exponentially with $1/d$ (see condition (C4)

in the Appendix). An indirect consequence of this is that the degree of the polynomials $P_i(\beta, x, p)$ and $P_i(\gamma, x, q)$ should increase when d increases (this follows from condition (C4)). For example, we will have to choose p and q at least equal to 2 when $d = 2$ (local quadratic estimators), and at least equal to 4 when $d = 3$ (order 4 local polynomial estimators).

Next, the test statistics are constructed from the difference between $F_{\varepsilon\theta_n}(y)$ and the nonparametric estimator of $F_\varepsilon(y)$:

$$\hat{F}_\varepsilon(y) = \frac{1}{n} \sum_{i=1}^n I(\hat{\varepsilon}_i \leq y). \quad (2.10)$$

This estimator was first studied by Akritas and Van Keilegom (2001) and then extended to the case where X is d -dimensional by Neumeyer and Van Keilegom (2008). Consider the process

$$W_n(y) = n^{1/2}(\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y)), \quad -\infty < y < \infty, \quad (2.11)$$

and define the following test statistics of the Kolmogorov-Smirnov and Cramér-von Mises types :

$$T_{KS} = n^{1/2} \sup_{-\infty < y < \infty} |\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y)|, \quad (2.12)$$

and

$$T_{CM} = n \int (\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y))^2 d\hat{F}_\varepsilon(y). \quad (2.13)$$

3 Asymptotic results

We now turn to the analysis of the asymptotic properties of the estimator θ_n and of the test statistics T_{KS} and T_{CM} . The assumptions under which these properties are valid, are given in the Appendix. The following notations will be used. Let $\dot{f}_{\varepsilon\theta}(y) = (\frac{\partial}{\partial\theta_1} f_{\varepsilon\theta}(y), \dots, \frac{\partial}{\partial\theta_k} f_{\varepsilon\theta}(y))^t$ and $f'_{\varepsilon\theta}(y) = \frac{d}{dy} f_{\varepsilon\theta}(y)$. Moreover, we will use the abbreviated notation $f_\varepsilon \equiv f_{\varepsilon\theta_0}$, $f'_\varepsilon \equiv f'_{\varepsilon\theta_0}$, $\dot{f}_\varepsilon \equiv \dot{f}_{\varepsilon\theta_0}$, and similarly for F_ε and \dot{F}_ε .

Theorem 3.1 *Assume (C1)-(C7). Then, under H_0 ,*

$$\theta_n - \theta_0 = -\Omega^{-1} n^{-1} \sum_{i=1}^n \xi(\varepsilon_i) + o_P(n^{-1/2}),$$

where

$$\xi(t) = \frac{\dot{f}_\varepsilon(t)}{f_\varepsilon(t)} + \int \frac{\dot{f}_\varepsilon(y)f'_\varepsilon(y)}{f_\varepsilon(y)} \left\{ t + \frac{y}{2}(t^2 - 1) \right\} dy,$$

and

$$\Omega = E \left[\frac{\dot{f}_\varepsilon(\varepsilon)\dot{f}_\varepsilon^t(\varepsilon)}{f_\varepsilon^2(\varepsilon)} \right].$$

Moreover,

$$n^{1/2}(\theta_n - \theta_0) \xrightarrow{d} N(0, \Omega^{-1}V\Omega^{-1}),$$

where $V = E[\xi(\varepsilon)\xi^t(\varepsilon)]$.

Theorem 3.2 *Assume (C1)-(C7). Then, under H_0 ,*

$$\begin{aligned} & \hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y) \\ &= n^{-1} \sum_{i=1}^n \left[I(\varepsilon_i \leq y) - F_\varepsilon(y) + \varphi(\varepsilon_i, y) + \dot{F}_\varepsilon^t(y) \Omega^{-1} \xi(\varepsilon_i) \right] + R_n(y), \end{aligned}$$

where $\sup_{-\infty < y < \infty} |R_n(y)| = o_P(n^{-1/2})$, and

$$\varphi(z, y) = f_\varepsilon(y) \left\{ z + \frac{y}{2}(z^2 - 1) \right\}.$$

Moreover, the process $n^{1/2}(\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y))$ ($-\infty < y < \infty$) converges weakly to a zero-mean Gaussian process $W(y)$ with covariance function

$$\begin{aligned} \text{Cov}(W(y_1), W(y_2)) &= E \left[\left\{ I(\varepsilon \leq y_1) - F_\varepsilon(y_1) + \varphi(\varepsilon, y_1) + \dot{F}_\varepsilon^t(y_1) \Omega^{-1} \xi(\varepsilon) \right\} \right. \\ &\quad \left. \times \left\{ I(\varepsilon \leq y_2) - F_\varepsilon(y_2) + \varphi(\varepsilon, y_2) + \dot{F}_\varepsilon^t(y_2) \Omega^{-1} \xi(\varepsilon) \right\} \right]. \end{aligned}$$

As a consequence of the above result, we now obtain the asymptotic limit of the test statistics T_{KS} and T_{CM} under H_0 .

Corollary 3.3 *Assume (C1)-(C7). Then, under H_0 ,*

$$T_{KS} \xrightarrow{d} \sup_{-\infty < y < \infty} |W(y)|,$$

and

$$T_{CM} \xrightarrow{d} \int W^2(y) dF_\varepsilon(y).$$

Remark 3.4 (Convergence under fixed alternatives) Note that if the error distribution F_ε is a fixed distribution (independent of the sample size n) that does not belong to the class \mathcal{F} , it can be easily seen that the test statistics T_{KS} and T_{CM} converge to

infinity. In fact, the estimators \hat{F}_ε and $F_{\varepsilon\theta_n}$ do not converge to the same distribution in that case, and hence the process $n^{1/2}(\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y))$, $-\infty < y < \infty$, diverges.

Remark 3.5 (Bootstrap approximation) To estimate the distributions of the statistics T_{KS} and T_{CM} under H_0 , the asymptotic result given in Corollary 3.3 could be used, with appropriate estimators for the unknown quantities. Alternatively, resampling techniques can provide very good precision. Here, the method we propose to use is as follows. For B fixed and for $b = 1, \dots, B$,

1. Let $\{\varepsilon_{1,b}^*, \dots, \varepsilon_{n,b}^*\}$ be an i.i.d. random sample from the distribution $F_{\varepsilon\theta_n}(\cdot)$.
2. Define new responses

$$Y_{i,b}^* = \hat{m}(X_i) + \hat{\sigma}(X_i)\varepsilon_{i,b}^*, \quad i = 1, \dots, n.$$

3. Let $T_{KS,b}^*$ and $T_{CM,b}^*$ be the test statistics obtained from the bootstrap sample $\{(X_1, Y_{1,b}^*), \dots, (X_n, Y_{n,b}^*)\}$.

Then, if we denote $T_{KS,(b)}^*$ for the b -th order statistic of $T_{KS,1}^*, \dots, T_{KS,B}^*$ and analogously for $T_{CM,(b)}^*$, then $T_{KS,((1-\alpha)B]+1}^*$ and $T_{CM,((1-\alpha)B]+1}^*$ approximate the $(1 - \alpha)$ -quantiles of the distributions of T_{KS} and T_{CM} respectively (where $[\cdot]$ denotes the integer part).

4 Practical implementation and simulations

In this section, we study the finite sample behavior of the proposed test statistics focusing on some main practical aspects of the implementation.

In the one-dimensional case ($d = 1$), we generate i.i.d. data from the model

$$Y = \alpha_0 + \alpha_1 X + \alpha_2 X^2 + \alpha_3 X^3 + (\beta_0 + \beta_1 X)\varepsilon, \quad (4.1)$$

where $X \sim U[0, 1]$, β_0 and β_1 are such that $\min(\beta_0, \beta_0 + \beta_1) > 0$, and ε is independent of X and is a mixture of two normal random variables, i.e. $F_\varepsilon(y) = p_1 P(Z_1 \leq y) + (1 - p_1)P(Z_2 \leq y)$, where Z_j is normal with mean μ_j and variance σ_j^2 ($j = 1, 2$). Clearly, the parameters $p_1, \mu_1, \sigma_1^2, \mu_2$ and σ_2^2 need to satisfy $p_1\mu_1 + (1 - p_1)\mu_2 = 0$ and $p_1(\sigma_1^2 + \mu_1^2) + (1 - p_1)(\sigma_2^2 + \mu_2^2) = 1$, since ε has zero mean and unit variance. Hence, $\mu_2 = -\mu_1 p_1 / (1 - p_1)$, and $\sigma_2^2 = \{\sigma_1^2 p_1^2 - (1 + \sigma_1^2 + \mu_1^2)p_1 + 1\} / (1 - p_1)^2$. We consider several values of p_1, μ_1 and σ_1^2 .

In the two-dimensional case, we generate i.i.d. data from the model

$$Y = \alpha_{10}X_1 + \alpha_{01}X_2 + \alpha_{11}X_1X_2 + \alpha_{20}X_1^2 + \alpha_{30}X_1^3 + (\beta_0 + \beta_{11}X_1X_2)\varepsilon, \quad (4.2)$$

where $X_1 \sim U[0, 1]$,

$$X_2|X_1 \sim \begin{cases} \beta(0.5, 0.5) & \text{if } X_1 \leq 0.1 \text{ or } X_1 \geq 0.9 \\ U[0, 1] & \text{if } 0.1 < X_1 < 0.9, \end{cases}$$

and for any $a, b > 0$, $\beta(a, b)$ is the beta-distribution with parameters a and b . The parameters β_0 and β_{11} satisfy $\min(\beta_0, \beta_{11}) > 0$, the error ε is independent of $X = (X_1, X_2)$ and has the same distribution as for model (4.1).

For each model, we simulate 500 samples of size $n = 100$ or 200 . For the estimation of $m(\cdot)$ and $\sigma(\cdot)$, we use the biweight kernel $k(u) = (15/16)(1 - u^2)^2I(-1 \leq u \leq 1)$ when $d = 1$ and the product kernel $k(u_1)k(u_2)$ when $d = 2$. The smoothing parameters h, l and g and the estimator θ_n are obtained from one of the six procedures described in Remark 2.1, namely by solving either (2.4), (2.5) and (2.3) (hereafter abbreviated by (a)), or (2.4), (2.6) and (2.3) (method (b)), or (2.7) and (2.8) for $s = 1$ (method (c)), or for $s = 2$ (method (d)), or (2.9) and (2.3) for $s = 1$ (method (e)), or for $s = 2$ (method (f)). When $d = 2$, both components of each bandwidth vector are chosen to be equal. The number of bootstrap replications is 200 and the level of the tests is 5%. Following Remark 2.2, we take $p = q = 1$, respectively $p = q = 2$ when X is one-dimensional, respectively two-dimensional.

We consider three different tests for F_ε :

1. $H_{01} : \varepsilon \sim \Phi(0, 1)$
2. $H_{02} : \varepsilon \sim p_1\Phi(0.9, 0.49) + (1 - p_1)\Phi\left(\frac{-0.9p_1}{1-p_1}, \frac{0.49p_1^2 - 2.3p_1 + 1}{(1-p_1)^2}\right)$
3. $H_{03} : \varepsilon \sim p_1\Phi(\mu_1, 0.49) + (1 - p_1)\Phi\left(\frac{-\mu_1p_1}{1-p_1}, \frac{0.49p_1^2 - (1.49 + \mu_1^2)p_1 + 1}{(1-p_1)^2}\right),$

where $\Phi(\mu, \sigma^2)$ stands for a normal distribution with mean μ and variance σ^2 . Clearly, hypothesis H_{01} is satisfied for $p_1 = 0, \mu_1 = 0$ and $\sigma_1 = 1$, hypothesis H_{02} is satisfied for any value of p_1 and for $\mu_1 = 0.9$ and $\sigma_1 = 0.7$, and hypothesis H_{03} is satisfied for any p_1 and μ_1 and for $\sigma_1 = 0.7$.

Tables 1 to 6 present the rejection proportions for the above three null hypotheses. In order to illustrate the results in a more complete way, we added a test based on a statistic

of the Anderson-Darling type given by

$$T_{AD} = n \int \frac{(\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y))^2}{\hat{F}_\varepsilon(y)(1 - \hat{F}_\varepsilon(y-))} d\hat{F}_\varepsilon(y). \quad (4.3)$$

A ‘parametric’ version of this statistic, obtained by replacing $d\hat{F}_\varepsilon(y)$ by $dF_{\varepsilon\theta_n}(y)$ and $\hat{F}_\varepsilon(y)(1 - \hat{F}_\varepsilon(y-))$ by $F_{\varepsilon\theta_n}(y)(1 - F_{\varepsilon\theta_n}(y-))$ in the expression above, was also studied but its results were slightly worse than those of T_{AD} . In the same way, the ‘parametric’ version of T_{CM} lead to quite similar results as T_{CM} . Rejection proportions for those ‘parametric’ statistics are therefore not reported here.

In general, it seems that rejection proportions under the alternatives are larger when (f) is used. However, under the null hypotheses, (c) seems to provide the closest results to 0.05, suggesting better bootstrap approximations of the distribution of the test statistics. In fact, the smoothing parameters obtained with (a), (b), (c) and (e) are usually smaller than those obtained with (d) and (f). In our cases, (a), (b), (c) and (e) often seem to overfit more or less the data such that the distributions of the corresponding test statistics seem more variable (suggesting less rejection under the alternatives). In some situations, this overfitting can also lead to more variability inside the resamples used by the bootstrap steps (also suggesting less rejection under the null). More precisely, beyond usual overfitting produced by the least squares cross validation procedures, two bandwidth parameters are involved in (a) and (b). In (a), g_{n1} is even obtained by adapting the data points $(Y_j - \hat{m}_{g,-j}(X_j))^2$, $j = 1, \dots, n$, to the kernel estimators $\hat{\sigma}_{g,-j}^2(X_j)$ through g in (2.5). As a consequence, we think (a) and (b) should only be used when strongly different behaviors of $m(\cdot)$ and $\sigma(\cdot)$ are detected. In (d) and (f), extrema of (2.7) and (2.9) are obtained for larger values of bandwidth parameters than in (c) and (e). Indeed, adding globally increasing values of $(Y_i - \hat{m}_{h,-i}(X_i))^2$ in the weighted average $\hat{\sigma}_{h,2}^2(X_j)$ (for increasing values of h) is more likely to increase the likelihood function, since it doesn’t consider the value of $(Y_j - \hat{m}_{h,-j}(X_j))^2$. Under the null, (c) seems to provide the closest resamples to the original samples, leading to good bootstrap approximations (and also suggesting that (c) is a good estimation procedure for the parameters of $F_{\varepsilon\theta}$). However, that doesn’t prevent large variabilities of the statistics obtained by (c) (and (e)) due to small h_{n1} (and h'_{n1}) and leading to small power. h_{n2} and h'_{n2} , that are globally larger, enable then to obtain better power since the corresponding statistics are less variable. Under H_1 , the best results are usually obtained by (f) since θ_n and h'_{n2} are obtained independently and without assuming H_0 .

When k increases (so when we move from H_{01} to H_{02} and H_{03}), the above consid-

p_1	μ_1	σ_1		(a)	(b)	(c)-(e)	(d)-(f)
0	0	1	T_{KS}	.046	.044	.040	.058
			T_{CM}	.038	.034	.042	.064
			T_{AD}	.038	0.36	.042	.062
0.4	0.9	0.7	T_{KS}	.232	.238	.226	.308
			T_{CM}	.284	.330	.288	.382
			T_{AD}	.242	.290	.266	.320
0.45	0.3	0.4	T_{KS}	.514	.550	.530	.548
			T_{CM}	.568	.598	.558	.652
			T_{AD}	.642	.690	.620	.756
0.44	0.9	0.7	T_{KS}	.630	.680	.652	.736
			T_{CM}	.740	.778	.756	.826
			T_{AD}	.708	.760	.738	.792
0.5	0.7	0.1	T_{KS}	1	1	1	1
			T_{CM}	1	1	1	1
			T_{AD}	1	1	1	1

Table 1: Rejection proportions under H_{01} for different mixtures of normal distributions and for model (4.1). The parameters determining $m(\cdot)$ and $\sigma(\cdot)$ are chosen as $\alpha_0 = 1$, $\alpha_1 = 1$, $\alpha_2 = -2$, $\alpha_3 = 1.5$, $\beta_0 = 0.1$ and $\beta_1 = 0.1$.

erations stay true, but obviously the bootstrap approximations are based on samples generated further and further from the true model and the distances between F_ε and \mathcal{F} become considerably smaller, inducing less and less power. However, as it turns out, (c) (resp. (f)) is less affected under H_0 (resp. H_1).

In Tables 4 to 6 we study the behavior of the different statistics for $d = 2$. Only the results for (f) are displayed. Indeed, the same characteristics as in the one-dimensional case apply. As can be expected, the power is smaller than for the one-dimensional case, but the results stay reasonably good. An increase of k seems to affect the results more than the same increase of d , and a small increase of n (from 100 to 200 in Table 6) already improves the results significantly.

p_1	μ_1	σ_1		(a)	(b)	(c)	(d)	(e)	(f)
0.44	0.9	0.7	T_{KS}	.044	.036	.038	.038	.046	.044
			T_{CM}	.038	.034	.046	.036	.044	.032
			T_{AD}	.042	.036	.050	.040	.046	.034
0	0	1	T_{KS}	.046	.042	.048	.054	.044	.056
			T_{CM}	.044	.054	.042	.056	.038	.062
			T_{AD}	.046	.048	.044	.052	.038	.056
0.45	0.3	0.4	T_{KS}	.756	.784	.686	.834	.682	.836
			T_{CM}	.788	.826	.732	.882	.734	.896
			T_{AD}	.788	.822	.718	.892	.724	.894
0.5	0.7	0.1	T_{KS}	1	1	1	1	1	1
			T_{CM}	1	1	1	1	1	1
			T_{AD}	1	1	1	1	1	1

Table 2: Rejection proportions under H_{02} for different mixtures of normal distributions and for model (4.1). The parameters determining $m(\cdot)$ and $\sigma(\cdot)$ are chosen as $\alpha_0 = 1$, $\alpha_1 = 1$, $\alpha_2 = 0$, $\alpha_3 = 0$, $\beta_0 = 0.3$ and $\beta_1 = 0.3$.

p_1	μ_1	σ_1		(a)	(b)	(c)	(d)	(e)	(f)
0.44	0.9	0.7	T_{KS}	.012	.034	.024	.010	.030	.018
			T_{CM}	.014	.018	.032	.016	.026	.010
			T_{AD}	.012	.016	.022	.016	.024	.010
0	0	1	T_{KS}	.022	.018	.028	.026	.018	.032
			T_{CM}	.016	.018	.024	.014	.020	.020
			T_{AD}	.016	.022	.018	.018	.030	.026
0.45	0.3	0.4	T_{KS}	.074	.080	.126	.136	.104	.164
			T_{CM}	.088	.116	.162	.192	.116	.210
			T_{AD}	.100	.136	.158	.206	.112	.252
0.5	0.7	0.1	T_{KS}	.978	.994	.998	1	.984	1
			T_{CM}	.962	.974	.996	1	.990	1
			T_{AD}	.974	.984	.996	1	.992	1

Table 3: Rejection proportions under H_{03} for different mixtures of normal distributions and for model (4.1). The parameters determining $m(\cdot)$ and $\sigma(\cdot)$ are chosen as $\alpha_0 = 1$, $\alpha_1 = 1$, $\alpha_2 = 0$, $\alpha_3 = 0$, $\beta_0 = 0.3$ and $\beta_1 = 0.3$.

p_1	μ_1	σ_1	T_{KS}	T_{CM}	T_{AD}
0	0	1	.062	.054	.074
0.4	0.9	0.7	.230	.244	.208
0.45	0.3	0.4	.432	.428	.446
0.44	0.9	0.7	.548	.634	.598
0.5	0.7	0.1	1	.996	1

Table 4: Rejection proportions under H_{01} for different mixtures of normal distributions and for model (4.2). The smoothing parameter is obtained from (f), $n = 100$, $\alpha_{10} = 1$, $\alpha_{01} = 1$, $\alpha_{11} = 1$, $\alpha_{20} = 4$, $\alpha_{30} = -3$, $\beta_0 = 0.3$ and $\beta_{11} = 0.3$.

p_1	μ_1	σ_1	T_{KS}	T_{CM}	T_{AD}
0.44	0.9	0.7	.032	.040	.038
0	0	1	.028	.022	.034
0.45	0.3	0.4	.448	.446	.454
0.5	0.7	0.1	1	.992	.998

Table 5: Rejection proportions under H_{02} for different mixtures of normal distributions and for model (4.2). The bandwidth parameter is obtained by (f), $n = 100$, $\alpha_{10} = 1$, $\alpha_{01} = 1$, $\alpha_{11} = 1$, $\alpha_{20} = 4$, $\alpha_{30} = -3$, $\beta_0 = 0.3$ and $\beta_{11} = 0.3$.

p_1	μ_1	σ_1	T_{KS}	T_{CM}	T_{AD}
0.45	0.3	0.4	.918	.934	.924

Table 6: Rejection proportions under H_{02} for one mixture of normal distributions and for model (4.2). The bandwidth parameter is obtained by (f), $n = 200$, $\alpha_{10} = 1$, $\alpha_{01} = 1$, $\alpha_{11} = 1$, $\alpha_{20} = 4$, $\alpha_{30} = -3$, $\beta_0 = 0.3$ and $\beta_{11} = 0.3$.

Appendix

In this appendix we state the assumptions under which the asymptotic results of Section 3 are valid, and we also give the proofs of these results. Throughout this appendix we will denote the true error by ε_0 , the true regression function by m_0 and the true scale function by σ_0 .

For an arbitrary θ , and for arbitrary functions m and $\sigma > 0$ defined on R_X , let

$$G(\theta, m, \sigma) = E \left[\frac{\dot{f}_{\varepsilon\theta} \left(\frac{Y-m(X)}{\sigma(X)} \right)}{f_{\varepsilon\theta} \left(\frac{Y-m(X)}{\sigma(X)} \right)} \right].$$

- (C1) For all $\delta > 0$, there exists an $\varepsilon > 0$ such that $\inf_{\|\theta-\theta_0\|>\delta} \|G(\theta, m_0, \sigma_0)\| > \varepsilon$.
- (C2) Uniformly for all $\theta \in \Theta$, $G(\theta, m, \sigma)$ is continuous with respect to the supremum norm in (m, σ) at $(m, \sigma) = (m_0, \sigma_0)$. Moreover, Ω is non-singular.
- (C3) k is a symmetric probability density function supported on $[-1, 1]$, k is d times continuously differentiable, and $k^{(j)}(\pm 1) = 0$ for $j = 0, \dots, d-1$.
- (C4) h_j, l_j and g_j are of the same order ($j = 1, \dots, d$) and satisfy $h_j/h^* \rightarrow b_j, l_j/h^* \rightarrow c_j$ and $g_j/h^* \rightarrow d_j$ for some $0 < b_j, c_j, d_j < \infty$ and some baseline bandwidth h^* . Moreover, for $r = p$ or q , h^* satisfies $nh^{*2r+4} \rightarrow 0$ when r is even, $nh^{*2r+2} \rightarrow 0$ when r is odd and $nh^{*3d+\delta} \rightarrow \infty$ for some small $\delta > 0$.
- (C5) All partial derivatives of F_X up to order $2d+1$ exist on the interior of R_X , they are uniformly continuous and $\inf_{x \in R_X} f_X(x) > 0$.
- (C6) All partial derivatives of m_0 and σ_0 up to order $p+2$ exist on the interior of R_X , they are uniformly continuous and $\inf_{x \in R_X} \sigma_0(x) > 0$.
- (C7) All (mixed) derivatives upto order 3 of $F_{\varepsilon\theta}(y)$ with respect to y and the components of θ exist and are continuous. Moreover, $\sup_y |y^2 f'_\varepsilon(y)| < \infty$ and $E(\varepsilon_0^6) < \infty$.

Proof of Theorem 3.1. For an arbitrary θ , and for arbitrary functions $m : R_X \rightarrow \mathbb{R}$ and $\sigma : R_X \rightarrow \mathbb{R}^+$, let

$$G_n(\theta, m, \sigma) = n^{-1} \sum_{i=1}^n \frac{\dot{f}_{\varepsilon\theta} \left(\frac{Y_i-m(X_i)}{\sigma(X_i)} \right)}{f_{\varepsilon\theta} \left(\frac{Y_i-m(X_i)}{\sigma(X_i)} \right)}.$$

Then, θ_n is the solution of $G_n(\theta, \hat{m}, \hat{\sigma}) = 0$. Also note that θ_0 satisfies $G(\theta_0, m_0, \sigma_0) = 0$. We will start by showing the consistency of θ_n . This will be done by checking the conditions of Theorem 1 in Chen, Linton and Van Keilegom (2003) (CLV hereafter). Condition (1.1) holds by definition of θ_n , while the second and third condition are guaranteed by assumptions (C1) and (C2). Condition (1.4) follows from Lemma A.1 in Neumeyer and Van Keilegom (2008), and from conditions (C3)-(C7). Finally, condition (1.5) is very similar to condition (2.5) of Theorem 2 of CLV, and we will verify both conditions below. So, the conditions of Theorem 1 are verified, up to condition (1.5) which we postpone to later. Next, we verify conditions (2.1)–(2.6) of Theorem 2 in CLV in order to obtain an i.i.d. representation for $\theta_n - \theta_0$ and to prove the asymptotic normality of θ_n . Condition (2.1) is, as for condition (1.1), valid by construction of the estimator θ_n . For condition (2.2), write for any $\theta \in \Theta$,

$$\begin{aligned}\Gamma_1(\theta) &:= \frac{\partial}{\partial \theta} G(\theta, m_0, \sigma_0) = E \left[\frac{\ddot{f}_{\varepsilon\theta}(\varepsilon_0) f_{\varepsilon\theta}(\varepsilon_0) - \dot{f}_{\varepsilon\theta}(\varepsilon_0) \dot{f}_{\varepsilon\theta}^t(\varepsilon_0)}{f_{\varepsilon\theta}^2(\varepsilon_0)} \right] \\ &= \int \ddot{f}_{\varepsilon\theta}(y) dy - \int \frac{\dot{f}_{\varepsilon\theta}(y) \dot{f}_{\varepsilon\theta}^t(y)}{f_{\varepsilon\theta}(y)} dy \\ &= \frac{\partial^2}{\partial \theta \partial \theta^t} \int f_{\varepsilon\theta}(y) dy - E \left[\frac{\dot{f}_{\varepsilon\theta}(\varepsilon_0) \dot{f}_{\varepsilon\theta}^t(\varepsilon_0)}{f_{\varepsilon\theta}^2(\varepsilon_0)} \right] = -E \left[\frac{\dot{f}_{\varepsilon\theta}(\varepsilon_0) \dot{f}_{\varepsilon\theta}^t(\varepsilon_0)}{f_{\varepsilon\theta}^2(\varepsilon_0)} \right],\end{aligned}$$

where $\ddot{f}_{\varepsilon\theta}(y) = \frac{\partial^2}{\partial \theta \partial \theta^t} f_{\varepsilon\theta}(y)$. Hence, condition (2.2) is easily seen to hold thanks to condition (C2) and (C7). As for condition (2.3), we need to calculate

$$\begin{aligned}\Gamma_2(\theta, m_0, \sigma_0)[m - m_0, \sigma - \sigma_0] \\ := \lim_{\tau \rightarrow 0} \frac{1}{\tau} \left\{ G(\theta, m_0 + \tau(m - m_0), \sigma_0 + \tau(\sigma - \sigma_0)) - G(\theta, m_0, \sigma_0) \right\}.\end{aligned}$$

For notational convenience, in a first stage, we ignore the σ component in the above expression and calculate

$$\begin{aligned}\Gamma_2(\theta, m_0)[m - m_0] &= \lim_{\tau \rightarrow 0} \frac{1}{\tau} \left\{ E \left[\frac{\dot{f}_{\varepsilon\theta}(Y - m_0(X) - \tau(m - m_0)(X))}{f_{\varepsilon\theta}(Y - m_0(X) - \tau(m - m_0)(X))} - \frac{\dot{f}_{\varepsilon\theta}(Y - m_0(X))}{f_{\varepsilon\theta}(Y - m_0(X))} \right] \right\} \\ &= -E \left[\frac{\dot{f}'_{\varepsilon\theta}(\varepsilon_0) f_{\varepsilon\theta}(\varepsilon_0) - \dot{f}_{\varepsilon\theta}(\varepsilon_0) f'_{\varepsilon\theta}(\varepsilon_0)}{f_{\varepsilon\theta}^2(\varepsilon_0)} (m - m_0)(X) \right] \\ &= - \left\{ \int \dot{f}'_{\varepsilon\theta}(y) dy - \int \frac{\dot{f}_{\varepsilon\theta}(y) f'_{\varepsilon\theta}(y)}{f_{\varepsilon\theta}(y)} dy \right\} E[(m - m_0)(X)] \\ &= \int \frac{\dot{f}_{\varepsilon\theta}(y) f'_{\varepsilon\theta}(y)}{f_{\varepsilon\theta}(y)} dy E[(m - m_0)(X)],\end{aligned}$$

since $\frac{\partial}{\partial \theta} \int f'_{\varepsilon\theta}(y) dy = 0$. Hence, taking the dependence on σ into account, we get that

$$\begin{aligned} & \Gamma_2(\theta, m_0, \sigma_0)[m - m_0, \sigma - \sigma_0] \\ &= \int \frac{\dot{f}_{\varepsilon\theta}(y) f'_{\varepsilon\theta}(y)}{f_{\varepsilon\theta}(y)} dy E\left[\frac{(m - m_0)(X)}{\sigma_0(X)}\right] + \int y \frac{\dot{f}_{\varepsilon\theta}(y) f'_{\varepsilon\theta}(y)}{f_{\varepsilon\theta}(y)} dy E\left[\frac{(\sigma - \sigma_0)(X)}{\sigma_0(X)}\right] \end{aligned}$$

for all θ , m and $\sigma > 0$. Now, it is easily seen that condition (2.3)(i) is satisfied, if we define for any function h defined on R_X ,

$$\|h\|_{d+\alpha} = \max_{k \leq d} \sup_{x \in R_X} |D^k h(x)| + \max_{k=d} \sup_{x, x' \in R_X} \frac{|D^k h(x) - D^k h(x')|}{\|x - x'\|^\alpha},$$

$$k = (k_1, \dots, k_d),$$

$$D^k = \frac{\partial^k}{\partial x_1^{k_1} \dots \partial x_d^{k_d}},$$

$k \cdot = \sum_{j=1}^d k_j$, and $\|\cdot\|$ is the Euclidean norm on \mathbb{R}^d . As for (2.3)(ii), note that it follows from the proof of Theorem 2 in CLV that it suffices to show that

$$\|\Gamma_2(\hat{\theta}, m_0, \sigma_0)[\hat{m} - m_0, \hat{\sigma} - \sigma_0] - \Gamma_2(\theta_0, m_0, \sigma_0)[\hat{m} - m_0, \hat{\sigma} - \sigma_0]\| \leq \|\hat{\theta} - \theta_0\|_{o_P(1)},$$

and this can be easily shown, using condition (C7). For condition (2.4), let

$$\mathcal{M} = \left\{ m : R_X \rightarrow \mathbb{R} : \|m\|_{d+\alpha} \leq M_1 \right\}$$

and

$$\mathcal{S} = \left\{ \sigma : R_X \rightarrow \mathbb{R} : \|\sigma\|_{d+\alpha} \leq M_1, \inf_{x \in R_X} \sigma(x) > M_0 \right\}$$

for some $0 < M_0 < M_1 < \infty$. Then, we apply once more (the proof of) Lemma A.1 in Neumeyer and Van Keilegom (2008) for the rate of convergence of \hat{m} and $\hat{\sigma}$. The same lemma ensures that $P(\hat{m} \in \mathcal{M}) \rightarrow 1$ and $P(\hat{\sigma} \in \mathcal{S}) \rightarrow 1$. Next, for verifying condition (2.5) we check the conditions of Theorem 3 in CLV. It suffices to check either condition (3.1) or (3.2). Condition (3.2) is verified for $s_j = 1$ and $j = 1, \dots, k$, whereas condition (3.3) follows from Theorem 2.7.1 in Van der Vaart and Wellner (1996). It remains to verify condition (2.6), which is immediate after applying the i.i.d. representation for $E\left[\frac{(\hat{m}-m_0)(X)}{\sigma_0(X)}\right]$ and $E\left[\frac{(\hat{\sigma}-\sigma_0)(X)}{\sigma_0(X)}\right]$, given in Lemma A.2 in Neumeyer and Van Keilegom (2008). This finishes the proof. \square

Proof of Theorem 3.2. Write

$$\begin{aligned} \hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y) &= [\hat{F}_\varepsilon(y) - F_\varepsilon(y)] - [F_{\varepsilon\theta_n}(y) - F_\varepsilon(y)] \\ &= n^{-1} \sum_{i=1}^n I(\varepsilon_i \leq y) - F_\varepsilon(y) + n^{-1} \sum_{i=1}^n \varphi(\varepsilon_i, y) \\ &\quad + \dot{F}_\varepsilon^t(y)(\theta_n - \theta_0) + o_P(n^{-1/2}), \end{aligned}$$

uniformly in y , where the latter equality follows from Theorem 2.1 in Neumeyer and Van Keilegom (2008). The result now follows from Theorem 3.1. \square

Proof of Corollary 3.3. The convergence of the Kolmogorov-Smirnov statistic T_{KS} follows directly from the continuous mapping theorem. For the Crámer-von Mises statistic T_{CM} it suffices to show that $d\hat{F}_\varepsilon(y)$ can be replaced by $dF_\varepsilon(y)$. Write

$$\begin{aligned} & \left| n \int (\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y))^2 d[\hat{F}_\varepsilon(y) - F_\varepsilon(y)] \right| \\ & \leq \left| n \int (\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y))^2 d[\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y)] \right| \\ & \quad + \left| n \int (\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y))^2 d[F_{\varepsilon\theta_n}(y) - F_\varepsilon(y)] \right|. \end{aligned}$$

It suffices to consider the second term above, which can be written as

$$\left| n \int (\hat{F}_\varepsilon(y) - F_{\varepsilon\theta_n}(y))^2 \dot{f}_{\varepsilon\tilde{\theta}}^t(y) dy \Omega^{-1} n^{-1} \sum_{i=1}^n \xi(\varepsilon_i) \right| + o_P(1) = o_P(1)$$

(with $\tilde{\theta}$ on the line segment between θ_0 and θ_n), which follows from Theorems 3.1 and 3.2. \square

References

- Akritis, M. G. and Van Keilegom, I. (2001). Nonparametric estimation of the residuals distribution. *Scand. J. Statist.*, **28**, 549–567.
- Alcalá, J.T., Cristóbal, J.A. and González Manteiga, W. (1999). Goodness-of-fit test for linear models based on local polynomials. *Statist. Probab. Letters*, **42**, 39–46.
- Chen, X., Linton, O. and Van Keilegom, I. (2003). Estimation of semiparametric models when the criterion function is not smooth. *Econometrica*, **71**, 1591–1608.
- Dette, H. (1999). A consistent test for the functional form of a regression based on a difference of variance estimators. *Ann. Statist.*, **27**, 1012–1040.
- Dette, H. and Munk, A. (1998). Validation of linear regression models. *Ann. Statist.*, **26**, 778–800.
- Dette, H., Neumeyer, N. and Van Keilegom, I. (2007). A new test for the parametric form of the variance function in nonparametric regression. *J. Royal Statist. Soc. - Series B*, **69**, 903–917.
- Fan, J., Zhang, C. and Zhang, J. (2001). Generalized likelihood ratio statistics and Wilks phenomenon. *Ann. Statist.*, **29**, 153–193.

- Härdle, W. and Mammen, E. (1993). Comparing non parametric versus parametric regression fits. *Ann. Statist.*, **21**, 1926–1947.
- Jiménez Gamero, M.D., Muñoz García, J. and Pino Mejías, R. (2005). Testing goodness of fit for the distribution of errors in multivariate linear models. *J. Multiv. Anal.*, **95**, 301–322.
- Neumeyer, N. Dette, H. and Nagel, E.-R. (2006). Bootstrap tests for the error distribution in linear and nonparametric regression models. *Aust. N. Z. J. Stat.*, **48**, 129-156.
- Neumeyer, N. and Van Keilegom, I. (2009). Estimating the error distribution in nonparametric multiple regression with applications to model testing. *J. Multiv. Anal.* (under revision).
- Stute, W. (1997). Nonparametric model checks for regression. *Ann. Statist.*, **25**, 613–641.
- Van der Vaart, A.W. and Wellner, J.A. (1996). *Weak convergence and empirical processes*. Springer, New York.
- Van Keilegom, I., González Manteiga, W. and Sánchez Sellero, C. (2008). Goodness-of-fit tests in parametric regression based on the estimation of the error distribution. *TEST*, **17**, 401–415.