

The INTEGER QUANTUM HALL EFFECT

Jean BELLISSARD

*Georgia Institute of Technology, Atlanta
School of Mathematics & School of Physics
e-mail: jeanbel@math.gatech.edu*

Sponsoring



Contributors

D. SPEHNER, *Institut Fourier*, Grenoble, France

H. SCHULZ-BALDES, *Dep. of Math.*, Friedrich-Alexander Universität, Erlangen-Nürnberg, Germany

A. VAN ELST

Main References

J. BELLISSARD, H. SCHULZ-BALDES, A. VAN ELST, "The Non Commutative Geometry of the Quantum Hall Effect"
J. Math. Phys., **35**, (1994), 5373-5471

A. CONNES, *Noncommutative Geometry*, Acad. Press, (1994)

J. E. AVRON, R. SEILER, B. SIMON, *Phys. Rev. Lett.*, **65**, (1990), 2185-2188.

J. E. AVRON, R. SEILER, B. SIMON, *Comm. Math. Phys.*, **159**, (1994), 399-422.

H. SCHULZ-BALDES, J. BELLISSARD, *Rev. Math. Phys.*, **10**, (1998), 1-46.

H. SCHULZ-BALDES, J. BELLISSARD, *J. Stat. Phys.*, **91**, (1998), 991-1026.


Content

1. Introduction to the QHE
2. Disorder and Magnetic Field
3. The Four Traces Way
4. Connes Formulae

I - INTRODUCTION to the IQHE

J. BELLISSARD, H. SCHULZ-BALDES, A. VAN ELST, *J. Math. Phys.*, **35**, (1994), 5373-5471

The Classical Hall Effect



B = magnetic field
 j = current density
 E = Hall electric field
 n = charge carrier density

In the stationary state:

$$en\vec{\mathcal{E}} + \vec{j} \times \vec{B} = 0$$

$$\Rightarrow \vec{j} = \begin{pmatrix} 0 & \sigma_H \\ -\sigma_H & 0 \end{pmatrix} \vec{\mathcal{E}}$$



$$\sigma_H = \frac{ne}{B}$$

$$\text{Units : } \frac{n}{B} = \left[\frac{1}{\text{flux}} \right], \quad \frac{h}{e} = [\text{flux}] \Rightarrow \nu = \frac{nh}{eB} = [1] = \textbf{(filling factor)}$$

This gives the *Hall formula*

$$\sigma_H = \frac{\nu}{R_H} \quad R_H = \frac{h}{e^2} = 25,812.80 \, \Omega$$


The Integer Quantum Hall Effect



Two examples of Hall bars used in experiments

(Gossard, 2000)

The Integer Quantum Hall Effect



J. P. EISENSTEIN, H. L. STORMER, *Science*, (1990), **248**, 1461

The Integer Quantum Hall Effect

- **Conditions of Observations**

- Low temperature ($\leq$ few Kelvins)
- Large sample size ($\geq$ few μm)
- High mobility & large quenched disorder
- Two-dimensional Fermion fluid

- **Experiment show that**

- Very flat plateaux at $\nu \sim 1, 2, 3, 4$ with $\sigma_H = \ell/R_H$, $\ell = 1, 2, 3, 4$
- Plateaux thickness $\delta\sigma_H/\sigma_H \leq 10^{-8} - 10^{-10}$
- Very small direct conductivity on plateaux $\Rightarrow$ *localization*
- For $\ell \geq 2$ electron-electron *interaction* is *negligible*

The Integer Quantum Hall Effect

- *Why is σ_H quantized ?*
- *What is the role of the localization ?*


Earlier Works: Laughlin's argument

R. B. LAUGHLIN, *Phys. Rev. B*, **23**, (1981), 5632

R. E. PRANGE, *Phys. Rev. B*, **23**, (1981), 4802

D. J. THOULESS, *J. Phys. C*, **14**, (1981), 3475


R. JOYNT, R. E. PRANGE, *Phys. Rev. B*, **29**, (1984), 3303



- Piercing the plane at x with a flux tube adiabatically varying from 0 to $\phi_0 = h/e$ forces one charge per filled Landau level to transfer from $x \rightarrow \infty$
- This adiabatic change induces a unitary transformation u on the Landau Hamiltonian (gauge transformation)
- This gives the quantization of the Hall conductance
- Localized states do not participate to this transport

Earlier Works: TKN₂

- Use the Harper model on a *square lattice*, nearest neighbor hopping terms, *uniform magnetic field* B perpendicular to the lattice
- Translation operators U_1, U_2



a = lattice spacing

ϕ = flux through unit cell

Earlier Works: TKN₂

- Commutation rules (*Rotation Algebra*)

$$U_1 U_2 = e^{2i\pi\alpha} U_2 U_1 \quad \alpha = \frac{\phi}{\phi_0} \quad \phi = Ba^2 \quad \phi_0 = \frac{h}{e}$$

- Kinetic Energy (*Hamiltonian*)

$$H = t(U_1 + U_2 + U_1^{-1} + U_2^{-1})$$

- Landau gauge $\psi(m, n) = e^{2i\pi mk} \varphi(n)$.

Hence $H\psi = E\psi$ means

$$\varphi(n+1) + \varphi(n-1) + 2 \cos 2\pi(n\alpha - k) \varphi(n) = \frac{E}{t} \varphi(n)$$

Earlier Works: TKN₂

- Choose $\alpha = p/q$ to make H q -periodic. Use *Bloch theory* with quasimomentum $\vec{k} = (k_1, k_2) \in \mathbb{B} \approx \mathbb{T}^2$
- At H is a $q \times q$ -matrix valued smooth function of $\vec{k}$
- At $\vec{k}$ fixed, any eigenstate $\Psi_{\vec{k}}$ of $H_{\vec{k}}$, defines a *line bundle* over $\mathbb{B}$
- Its non triviality is controlled by the *Chern number*

$$\text{Ch}(\Psi) = \frac{1}{\pi} \int_0^{2\pi} \int_0^{2\pi} \Im m \left\langle \frac{\partial \Psi}{\partial k_1} \middle| \frac{\partial \Psi}{\partial k_2} \right\rangle dk_1 dk_2$$

- $\text{Ch}(\Psi) \in \mathbb{Z}$ and is *homotopy invariant* under deformation of H

Earlier Works: TKN₂

- If $P : \vec{k} \in \mathbb{B} \mapsto P(\vec{k})$ is a *projection* valued smooth map then
(example: $P = |\Psi\rangle\langle\Psi|$)

$$\text{Ch}(P) = \frac{1}{2i\pi} \int_0^{2\pi} \int_0^{2\pi} \text{Tr} \left(P(\vec{k}) \left[\frac{\partial P}{\partial k_1}, \frac{\partial P}{\partial k_2} \right] \right) dk_1 dk_2 \in \mathbb{Z}$$

- If P, Q are two orthogonal projections, $PQ = QP = 0$, then

$$\text{Ch}(P \oplus Q) = \text{Ch}(P) + \text{Ch}(Q)$$

Earlier Works: TKN₂

- If the *Fermi level* E_F belongs to an energy gap, let P_F be the *Fermi projection* (namely the eigenprojection onto states with energy $E \leq E_F$)
- Then the following *Chinese-Japanese* relation holds

$$\sigma_H = \frac{e^2}{h} \mathbf{Ch}(P_F) \quad (\text{Chern-Kubo formula})$$

- This formula *explains the quantization* of the Hall conductance *for rational magnetic fields !*


It does NOT explain the appearance of plateaux !

II - Disorder and Magnetic Field


Noncommutativity of the Brillouin Zone

- If $\alpha = \phi/\phi_0 \notin \mathbb{Q}$, the Bloch theory *fails* !
- Adding a *random potential* adds up to the *failure* of Bloch theory !
- **Disordered potential:** $V_\omega(x) = W \omega_x, x \in \mathbb{Z}^2$ with
 - W is the disorder strength
 - $\omega = (\omega_x)_{x \in \mathbb{Z}^2}$ and the ω_x 's are *i.i.d.*'s with *uniform distribution* on $[-1/2, +1/2]$
 - $\omega \in \Omega = \prod_{x \in \mathbb{Z}^2} [-1/2, +1/2]$ is compact (*Tychonov Theorem*) and $\mathbb{Z}^2$ acts by *shift*.
- The groupoid is now $\Omega \rtimes \mathbb{Z}^2$
The observable algebra is again $\mathcal{A} = C(\Omega) \rtimes_B \mathbb{Z}^d$.

Landau Levels



Landau Levels




Landau levels

$$E_\ell = \ell\alpha$$


$$\ell = 1, 2, \dots$$

Density of States



$$\int_{\mathbb{R}} f(E) g(E) dE = \mathcal{T}_{\mathbb{P}}(f(H))$$

(for $f \in C_c(\mathbb{R})$)



NO gap !

Noncommutativity of the Brillouin Zone

- The *Chern-Kubo formula* becomes

$$\sigma_H = -2i\pi \frac{e^2}{h} \mathcal{T}_{\mathbb{P}}(P_F [\partial_1 P_F, \partial_2 P_F]) = \frac{e^2}{h} \mathbf{Ch}(P_F)$$

- **Questions:**

- How does one prove that $\mathbf{Ch}(P_F) \in \mathbb{Z}$?
- How does one define $\mathbf{Ch}(P_F)$ if the Fermi level *does NOT belong* to a gap !

III - The Four Traces Way

J. BELLISSARD, H. SCHULZ-BALDES, A. VAN ELST, *J. Math. Phys.*, **35**, (1994), 5373-5471

Trace and Trace per Unit Volume

- For a *trace class* operator T acting on a separable Hilbert space

$$\mathrm{Tr}(T) = \sum_{n=1}^{\infty} \langle e_n | T e_n \rangle \quad (e_n)_{n \in \mathbb{N}} \text{ orthonormal basis}$$

- If Γ is a locally compact *groupoid*, with unit space $\mathbb{E}$ equipped with an invariant *probability* measure $\mathbb{P}$

$$\mathcal{T}_{\mathbb{P}}(A) = \int_{\mathbb{E}} A(\xi, 0) d\mathbb{P}(\xi) \quad A \in C_c(\Gamma)$$

Graded Trace

- **Spinors:** here $\Gamma^\xi \subset \mathbb{R}^2$! If $\mathcal{H}_\xi = L^2(\Gamma^\xi)$ set $\widehat{H}_\xi = H_\xi \otimes \mathbb{C}^2$
- **Grading:**

$$G = \mathbf{1}_{\mathcal{H}_\xi} \otimes \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad G^* = G = G^{-1}$$

- An operator $T \in \mathcal{B}(\mathcal{H}_\xi)$ has *degree* $\deg(T)$ whenever

$$GT - (-1)^{\deg(T)} TG = 0$$

- Any operator $T \in \mathcal{B}(\mathcal{H}_\xi)$ can be uniquely decomposed into $T = T_0 + T_1$ with $\deg(T_i) = i$

Graded Trace

- **Graded Commutator:**

$$[T, T']_S = TT' - (-1)^{\deg(T) \deg(T')} T'T$$

- **Dirac Operator:** if $X = X_1 + \iota X_2$ is the *position* operator

$$D = \begin{bmatrix} 0 & X \\ X^* & 0 \end{bmatrix} \quad F = \frac{D}{|D|} \Rightarrow F = F^* = F^{-1}, \deg(F) = 1$$

- **Graded Trace:**

$$\mathrm{Tr}_S(T) = \frac{1}{2} \mathrm{Tr} (GF [F, T]_S)$$

Graded Trace

- **Differential:**

$$dT = [F, T]_S$$

- **Leibniz rule:**

$$d(TT') = dT T' + (-1)^{\deg(T)} T dT'$$

- Tr_S is *linear* and satisfies, for dT, dT' *trace class* operator

$$\text{Tr}_S(TT') = (-1)^{\deg(T) \deg(T')} \text{Tr}_S(T'T) \quad \text{(graded trace)}$$

Graded Trace

- **Representation of $C^*(\Gamma)$**

$$\widehat{\pi}_\xi(A) = \begin{bmatrix} \pi_\xi(A) & 0 \\ 0 & \pi_\xi(A) \end{bmatrix} \quad A \in \mathcal{A}_0, \deg(\widehat{\pi}_\xi(A)) = 0$$

- **Laughlin argument:** It is worth noticing that $u = X/|X|$ is a unitary operator on $\mathcal{H}_\xi$ representing the *gauge transformation* corresponding to an *adiabatic change* of a pointwise flux at the origin, from 0 to ϕ_0 .

The Dixmier Trace

J. DIXMIER, *C. R. Acad. Sci. Paris Sér. A-B*, **262**, (1966), A1107-A1108

- If $\mathcal{H}$ is a Hilbert space $L^p(\mathcal{H})$ denotes the *Schatten ideal* of compact operators with $\text{Tr}(|T|^p) < \infty$
- If T is compact, let $\mu_1 \geq \cdots \geq \mu_n \geq 0$ be its singular values (eigenvalues of $|T|$) labelled in nonincreasing order. Then

$$\|T\|_{p+} = \left(\limsup_{N \in \mathbb{N}} \frac{1}{\ln(N+1)} \sum_{n=1}^N \mu_n^p \right)^{1/p}$$

- **Mačaev ideal:** $L^{p+}(\mathcal{H})$ is the set of T compact with $\|T\|_{p+} < \infty$

The Dixmier Trace

Theorem *Let $L^{p-}(\mathcal{H}) = \{T \text{ compact} ; \|T\|_{p+} = 0\}$. Then*

1. $L^{p\pm}(\mathcal{H})$ are two-sided ideals in $\mathcal{B}(\mathcal{H})$

2. If $0 \leq p < p' < \infty$

$$L^p(\mathcal{H}) \subset L^{p-}(\mathcal{H}) \subset L^{p+}(\mathcal{H}) \subset L^{p'}(\mathcal{H})$$

3. $\|\cdot\|_{p+}$ is a seminorm making $L^{p+}(\mathcal{H})/L^{p-}(\mathcal{H})$ a Banach space

The Dixmier Trace

- **Abstract nonsense:** using the theory of amenable groups, Dixmier proves the existence of a *linear form* $\Upsilon : \ell^\infty(\mathbb{N}) \rightarrow \mathbb{R}$ such that
 - $\Upsilon(a_1, a_2, a_3, \dots) = \Upsilon(a_2, a_3, a_4, \dots)$
 - $\Upsilon(a_1, a_2, a_3, \dots) = \Upsilon(a_1, a_1, a_2, a_2, \dots)$
 - If $a \in \ell^\infty(\mathbb{N})$ converges, then $\Upsilon(a) = \lim_{n \rightarrow \infty} (a_n)$
- **Dixmier Trace:** given such Υ , then

$$\mathrm{Tr}_\Upsilon(T) = \Upsilon \left(\frac{1}{\ln(N+1)} \sum_{n=1}^N \mu \right) \quad T \in L^{1+}(\mathcal{H}), \quad T \geq 0$$

The Dixmier Trace

- Then Dixmier proves that Tr_γ extends as a *positive linear map* on $L^{p^+}(\mathcal{H})$ vanishing on $L^{p^-}(\mathcal{H})$ and such that

$$\text{Tr}_\gamma(UTU^{-1}) = \text{Tr}_\gamma(T)$$

$$\text{Tr}_\gamma(ST) = \text{Tr}_\gamma(TS)$$

if U is unitary and $S, T \in L^{p^+}(\mathcal{H})$

IV - Connes Formulae

A. CONNES, *Noncommutative Geometry*, Acad. Press, (1994)

First Connes Formula

- If $A \in C^*(\Gamma)$ then for $\mathbb{P}$ -almost all ξ 's and all Υ

$$\mathcal{T}_{\mathbb{P}}(|\vec{\nabla}A|^2) \stackrel{def}{=} \mathcal{T}_{\mathbb{P}}(|\partial_1 A|^2 + |\partial_2 A|^2) = \frac{1}{\pi} \operatorname{Tr}_{\Upsilon}(|d\pi_{\xi}(A)|^2)$$

- If $\mathcal{S}$ denotes the *Sobolev space* generated by $A \in \mathcal{A}_0$ such that $\mathcal{T}_{\mathbb{P}}(|A|^2 + |\vec{\nabla}A|^2) < \infty$ then

$$A \in \mathcal{S} \Rightarrow d\pi_{\xi}(A) \in L^{2+}(\mathcal{H})$$

Second Connes Formula

- **A cyclic 2-cocycle:** for $A_0, A_1, A_2 \in \mathcal{S}$

$$\mathcal{T}_2(A_0, A_1, A_2) = 2i\pi \mathcal{T}_{\mathbb{P}}(A_0 (\partial_1 A_1 \partial_2 A_2 - \partial_2 A_1 \partial_1 A_2))$$

- **Cyclicity:**

$$\mathcal{T}_2(A_0, A_1, A_2) = \mathcal{T}_2(A_2, A_0, A_1)$$

- **$\mathcal{T}_2$ is Hochschild-closed:**

$$\begin{aligned} (b\mathcal{T}_2)(A_0, A_1, A_2, A_3) &= \mathcal{T}_2(A_0 A_1, A_2, A_3) - \mathcal{T}_2(A_0, A_1 A_2, A_3) \\ &\quad + \mathcal{T}_2(A_0, A_1, A_2 A_3) - \mathcal{T}_2(A_3 A_0, A_1, A_2) \\ &= 0 \end{aligned}$$

Second Connes Formula

- for $A_0, A_1, A_2 \in \mathcal{S}$

$$\mathcal{T}_2(A_0, A_1, A_2) = \int_{\Xi} \text{Tr}_{\mathcal{S}}(\widehat{\pi}_{\xi}(A_0) \, d\widehat{\pi}_{\xi}(A_1) \, d\widehat{\pi}_{\xi}(A_2)) \, d\mathbb{P}(\xi)$$

•

$$\text{Tr}_{\mathcal{S}}(\widehat{\pi}_{\xi}(A_0) \, d\widehat{\pi}_{\xi}(A_1) \, d\widehat{\pi}_{\xi}(A_2)) = \frac{1}{2} \text{Tr}(G \, d\widehat{\pi}_{\xi}(A_0) \, d\widehat{\pi}_{\xi}(A_1) \, d\widehat{\pi}_{\xi}(A_2))$$

- $A_i \in \mathcal{S} \Rightarrow d\widehat{\pi}_{\xi}(A) \in L^{2+}(\mathcal{H}) \subset L^3(\mathcal{H})$ so that the *r.h.s* is well defined

Fredholm Index

- **Integrality:** (Connes) If P is a projection on $\mathcal{H}_\xi$ then set $\widehat{P} = P \otimes \mathbf{1}_2$. If $d\widehat{P} \in L^3(\mathcal{H})$ then PuP is *Fredholm* and

$$\mathrm{Tr}_S(\widehat{P} d\widehat{P} d\widehat{P}) = \mathrm{Ind}(PuP) \in \mathbb{Z}$$

- **Fedosov formula:** $d\widehat{P} \in L^3(\mathcal{H}) \Leftrightarrow (PuP - P) \in L^3(\mathcal{H})$ and

$$\mathrm{Ind}(PuP) = \mathrm{Tr}((PuP - P)^{2n+1}) \quad \forall n \geq 1$$

- Hence $\mathrm{Ind}(PuP)$ measure the *change of dimension* of P under the Laughlin gauge transformation, namely the *number of charges* sent to infinity (Avron, Seiler, Simon)

Quantization of the Chern Number

- If $P_F \in \mathcal{S}$, then $d\widehat{\pi}_\xi(P_F) \in L^{2+}(\mathcal{H}_\xi) \subset L^3(\mathcal{H}_\xi)$ (1st Connes formula)
- Then (2nd Connes formula)

$$\mathbf{Ch}(P_F) = \int_{\Xi} \mathrm{Tr}_{\mathcal{S}}(\widehat{\pi}_\xi(P_F) d\widehat{\pi}_\xi(P_F) \widehat{\pi}_\xi(P_F)) d\mathbb{P}(\xi) = \int_{\Xi} n(\xi) d\mathbb{P}(\xi)$$

- Since it is a Fredholm index $n(\xi) \in \mathbb{Z}$. By covariance it is *translation invariant*. Since $P_F \in \mathcal{S}$ it follows that $n(\xi)$ is *measurable* in ξ . Since $\mathbb{P}$ is *ergodic*, then $n(\xi)$ is almost surely constant. Hence

$$P_F \in \mathcal{S} \Rightarrow \mathbf{Ch}(P_F) \in \mathbb{Z}$$

which measures the *number of charges* sent to infinity.

Localization

The condition $P_F \in \mathcal{S}$ is implied by the condition that the Fermi level E_F lies in a region of *localized states*

Thanks for listening !